

So what? | Lagrangians
to
Lasers



Search for new physics at the Large Hadron Collider

...

SHREYAS BAKARE

The background of the slide is a dark, blue-grey image of the Large Hadron Collider (LHC) tunnel. The tunnel is a long, cylindrical structure that recedes into the distance, creating a strong sense of perspective. The lighting is dim, with some highlights on the tunnel's surface, emphasizing its scale and depth.

[?]Search [?]for new physics at the _?Large Hadron Collider

...

SHREYAS BAKARE




Search for new physics at the Large Hadron Collider

...


SHREYAS BAKARE

*Experimental High Energy
Physics*


Energy scales in physics



High Energy Physics



Standard Model




$$\begin{aligned}
 & -\frac{1}{2}\partial_\nu g_\mu^a \partial_\nu g_\mu^a - g_s f^{abc} \partial_\mu g_\nu^a g_\mu^b g_\nu^c - \frac{1}{4} g_s^2 f^{abc} f^{ade} g_\mu^b g_\nu^c g_\mu^d g_\nu^e + \\
 & \frac{1}{2} i g_s^2 (\bar{q}_i^\mu \gamma^\mu q_j^\mu) g_\mu^a + \bar{G}^a \partial^2 G^a + g_s f^{abc} \partial_\mu \bar{G}^a G^b g_\mu^c - \partial_\nu W_\mu^+ \partial_\nu W_\mu^- - \\
 & M^2 W_\mu^+ W_\mu^- - \frac{1}{2} \partial_\nu Z_\mu^0 \partial_\nu Z_\mu^0 - \frac{1}{2 c_w^2} M^2 Z_\mu^0 Z_\mu^0 - \frac{1}{2} \partial_\mu A_\nu \partial_\mu A_\nu - \frac{1}{2} \partial_\mu H \partial_\mu H - \\
 & \frac{1}{2} m_h^2 H^2 - \partial_\mu \phi^+ \partial_\mu \phi^- - M^2 \phi^+ \phi^- - \frac{1}{2} \partial_\mu \phi^0 \partial_\mu \phi^0 - \frac{1}{2 c_w^2} M \phi^0 \phi^0 - \beta_h \left[\frac{2 M^2}{g^2} + \right. \\
 & \left. \frac{2 M}{g} H + \frac{1}{2} (H^2 + \phi^0 \phi^0 + 2 \phi^+ \phi^-) \right] + \frac{2 M^4}{g^2} \alpha_h - i g c_w [\partial_\nu Z_\mu^0 (W_\mu^+ W_\nu^- - \\
 & W_\nu^+ W_\mu^-) - Z_\nu^0 (W_\mu^+ \partial_\nu W_\mu^- - W_\nu^- \partial_\mu W_\mu^+) + Z_\mu^0 (W_\nu^+ \partial_\nu W_\mu^- - \\
 & W_\nu^- \partial_\mu W_\mu^+) - i g s_w [\partial_\nu A_\mu (\bar{W}_\mu^+ W_\nu^- - W_\nu^+ \bar{W}_\mu^-) - A_\nu (W_\mu^+ \partial_\nu \bar{W}_\mu^- - \\
 & W_\mu^- \partial_\nu \bar{W}_\mu^+) + A_\mu (W_\nu^+ \partial_\nu \bar{W}_\mu^- - W_\nu^- \partial_\mu \bar{W}_\mu^+)] - \frac{1}{2} g^2 W_\mu^+ W_\mu^- W_\nu^+ W_\nu^- + \\
 & \frac{1}{2} g^2 W_\mu^+ W_\nu^- W_\mu^+ W_\nu^- + g^2 c_w^2 (Z_\mu^0 W_\mu^+ Z_\nu^0 W_\nu^- - Z_\mu^0 Z_\nu^0 W_\mu^+ W_\nu^-) + \\
 & g^2 s_w^2 (A_\mu W_\mu^+ A_\nu W_\nu^- - A_\mu A_\nu W_\mu^+ W_\nu^-) + g^2 s_w c_w [A_\mu Z_\nu^0 (W_\mu^+ W_\nu^- - \\
 & W_\nu^+ W_\mu^-) - 2 A_\mu Z_\mu^0 W_\nu^+ W_\nu^-] - g \alpha [H^3 + H \phi^0 \phi^0 + 2 H \phi^+ \phi^-] - \\
 & \frac{1}{8} g^2 \alpha_h [H^4 + (\phi^0)^4 + 4 (\phi^+ \phi^-)^2 + 4 (\phi^0)^2 \phi^+ \phi^- + 4 H^2 \phi^+ \phi^- + 2 (\phi^0)^2 H^2] - \\
 & g M W_\mu^+ W_\mu^- H - \frac{1}{2} g \frac{M}{c_w^2} Z_\mu^0 Z_\mu^0 H - \frac{1}{2} i g [W_\mu^+ (\phi^0 \partial_\mu \phi^- - \phi^- \partial_\mu \phi^0) - \\
 & W_\mu^- (\phi^0 \partial_\mu \phi^+ - \phi^+ \partial_\mu \phi^0)] + \frac{1}{2} i g [W_\mu^+ (H \partial_\mu \phi^- - \phi^- \partial_\mu H) - W_\mu^- (H \partial_\mu \phi^+ - \\
 & \phi^+ \partial_\mu H)] + \frac{1}{2} g \frac{1}{c_w} (Z_\mu^0 (H \partial_\mu \phi^0 - \phi^0 \partial_\mu H) - i g \frac{s_w^2}{c_w} M Z_\mu^0 (W_\mu^+ \phi^- - W_\mu^- \phi^+) + \\
 & i g s_w M A_\mu (W_\mu^+ \phi^- - W_\mu^- \phi^+) - i g \frac{1 - 2 c_w^2}{2 c_w} Z_\mu^0 (\phi^+ \partial_\mu \phi^- - \phi^- \partial_\mu \phi^+) + \\
 & i g s_w A_\mu (\phi^+ \partial_\mu \phi^- - \phi^- \partial_\mu \phi^+) - \frac{1}{4} g^2 W_\mu^+ W_\mu^- [H^2 + (\phi^0)^2 + 2 \phi^+ \phi^-] - \\
 & \frac{1}{4} g^2 \frac{1}{c_w^2} Z_\mu^0 Z_\mu^0 [H^2 + (\phi^0)^2 + 2 (2 s_w^2 - 1) \phi^+ \phi^-] - \frac{1}{2} g^2 \frac{s_w^2}{c_w^2} Z_\mu^0 \phi^0 (W_\mu^+ \phi^- + \\
 & W_\mu^- \phi^+) - \frac{1}{2} i g^2 \frac{s_w^2}{c_w} Z_\mu^0 H (W_\mu^+ \phi^- - W_\mu^- \phi^+) + \frac{1}{2} g^2 s_w A_\mu \phi^0 (W_\mu^+ \phi^- + \\
 & W_\mu^- \phi^+) + \frac{1}{2} i g^2 s_w A_\mu H (W_\mu^+ \phi^- - W_\mu^- \phi^+) - g^2 \frac{s_w}{c_w} (2 c_w^2 - 1) Z_\mu^0 A_\mu \phi^+ \phi^- - \\
 & g^4 s_w^2 A_\mu A_\nu \phi^+ \phi^- - \bar{e}^\lambda (\gamma^\mu \partial + m_e^\lambda) e^\lambda - \bar{\nu}^\lambda \gamma^\mu \partial \nu^\lambda - \bar{u}_j^\lambda (\gamma^\mu \partial + m_u^\lambda) u_j^\lambda - \\
 & \bar{d}_j^\lambda (\gamma^\mu \partial + m_d^\lambda) d_j^\lambda + i g s_w A_\mu [(-\bar{e}^\lambda \gamma^\mu e^\lambda) + \frac{2}{3} (\bar{u}_j^\lambda \gamma^\mu u_j^\lambda) - \frac{1}{3} (\bar{d}_j^\lambda \gamma^\mu d_j^\lambda)] + \\
 & \frac{i g}{4 c_w} Z_\mu^0 [(\bar{\nu}^\lambda \gamma^\mu (1 + \gamma^5) \nu^\lambda) + (\bar{e}^\lambda \gamma^\mu (4 s_w^2 - 1 - \gamma^5) e^\lambda) + (\bar{u}_j^\lambda \gamma^\mu (\frac{4}{3} s_w^2 - \\
 & 1 - \gamma^5) u_j^\lambda) + (\bar{d}_j^\lambda \gamma^\mu (1 - \frac{8}{3} s_w^2 - \gamma^5) d_j^\lambda)] + \frac{i g}{2 \sqrt{2}} W_\mu^+ [(\bar{\nu}^\lambda \gamma^\mu (1 + \gamma^5) e^\lambda) + \\
 & (\bar{u}_j^\lambda \gamma^\mu (1 + \gamma^5) C_{\lambda k} d_j^k)] + \frac{i g}{2 \sqrt{2}} W_\mu^- [(\bar{e}^\lambda \gamma^\mu (1 + \gamma^5) \nu^\lambda) + (\bar{d}_j^\lambda C_{\lambda k}^\dagger \gamma^\mu (1 + \\
 & \gamma^5) u_j^k)] + \frac{i g}{2 \sqrt{2}} \frac{m_\lambda^2}{M} [-\phi^+ (\bar{\nu}^\lambda (1 - \gamma^5) e^\lambda) + \phi^- (\bar{e}^\lambda (1 + \gamma^5) \nu^\lambda)] - \\
 & \frac{g}{2} \frac{m_\lambda^2}{M} [H (\bar{e}^\lambda e^\lambda) + i \phi^0 (\bar{e}^\lambda \gamma^5 e^\lambda)] + \frac{i g}{2 M \sqrt{2}} \phi^+ [-m_d^\lambda (\bar{u}_j^\lambda C_{\lambda k} (1 - \gamma^5) d_j^k) + \\
 & m_u^\lambda (\bar{u}_j^\lambda C_{\lambda k} (1 + \gamma^5) d_j^k) + \frac{i g}{2 M \sqrt{2}} \phi^- [m_d^\lambda (\bar{d}_j^\lambda C_{\lambda k}^\dagger (1 + \gamma^5) u_j^k) - m_u^\lambda (\bar{d}_j^\lambda C_{\lambda k}^\dagger (1 - \\
 & \gamma^5) u_j^k) - \frac{g}{2} \frac{m_\lambda^2}{M} H (\bar{u}_j^\lambda u_j^\lambda) - \frac{g}{2} \frac{m_\lambda^2}{M} H (\bar{d}_j^\lambda d_j^\lambda) + \frac{i g}{2} \frac{m_\lambda^2}{M} \phi^0 (\bar{u}_j^\lambda \gamma^5 u_j^\lambda) - \\
 & \frac{i g}{2} \frac{m_\lambda^2}{M} \phi^0 (\bar{d}_j^\lambda \gamma^5 d_j^\lambda) + \bar{X}^+ (\partial^2 - M^2) X^+ + \bar{X}^- (\partial^2 - M^2) X^- + \bar{X}^0 (\partial^2 - \\
 & \frac{M^2}{c_w^2}) X^0 + \bar{Y} \partial^2 Y + i g c_w W_\mu^+ (\partial_\mu \bar{X}^0 X^- - \partial_\mu \bar{X}^+ X^0) + i g s_w W_\mu^+ (\partial_\mu \bar{Y} X^- - \\
 & \partial_\mu \bar{X}^+ Y) + i g c_w W_\mu^- (\partial_\mu \bar{X}^- X^0 - \partial_\mu \bar{X}^0 X^+) + i g s_w W_\mu^- (\partial_\mu \bar{X}^- Y - \\
 & \partial_\mu \bar{Y} X^+) + i g c_w Z_\mu^0 (\partial_\mu \bar{X}^+ X^+ - \partial_\mu \bar{X}^- X^-) + i g s_w A_\mu (\partial_\mu \bar{X}^+ X^+ - \\
 & \partial_\mu \bar{X}^- X^-) - \frac{1}{2} g M [\bar{X}^+ X^+ H + \bar{X}^- X^- H + \frac{1}{c_w} \bar{X}^0 X^0 H] +
 \end{aligned}$$



Standard Model

Three generations of matter (fermions)

	I	II	III	
mass →	2.4 MeV/c ²	1.27 GeV/c ²	171.2 GeV/c ²	0
charge →	2/3	2/3	2/3	0
spin →	1/2	1/2	1/2	1
name →	u up	c charm	t top	γ photon
Quarks	4.8 MeV/c ²	104 MeV/c ²	4.2 GeV/c ²	0
	-1/3	-1/3	-1/3	0
	1/2	1/2	1/2	1
	d down	s strange	b bottom	g gluon
Leptons	<2.2 eV/c ²	<0.17 MeV/c ²	<15.5 MeV/c ²	91.2 GeV/c ²
	0	0	0	0
	1/2	1/2	1/2	1
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	Z⁰ Z boson
Gauge bosons	0.511 MeV/c ²	105.7 MeV/c ²	1.777 GeV/c ²	80.4 GeV/c ²
	-1	-1	-1	±1
	1/2	1/2	1/2	1
	e electron	μ muon	τ tau	W[±] W boson




$$\begin{aligned}
 & -\frac{1}{2}\partial_\nu g_\mu^\nu g_\mu^\mu - g_s f^{abc}\partial_\mu g_\nu^a g_\mu^b g_\nu^c - \frac{1}{4}g_s^2 f^{abc}f^{ade}g_\mu^b g_\nu^c g_\mu^d g_\nu^e + \\
 & \frac{1}{2}g_s^2(\bar{q}_i^\gamma \gamma^\mu q_j^\gamma)g_\mu^\gamma + \bar{G}^a\partial^2 G^a + g_s f^{abc}\partial_\mu \bar{G}^a G^b g_\mu^c - \partial_\nu W_\mu^+ \partial_\nu W_\mu^- - \\
 & M^2 W_\mu^+ W_\mu^- - \frac{1}{2}\partial_\nu Z_\mu^0 \partial_\nu Z_\mu^0 - \frac{1}{2c_w^2}M^2 Z_\mu^0 Z_\mu^0 - \frac{1}{2}\partial_\mu A_\nu \partial_\mu A_\nu - \frac{1}{2}\partial_\mu H \partial_\mu H - \\
 & \frac{1}{2}m_h^2 H^2 - \partial_\mu \phi^+ \partial_\mu \phi^- - M^2 \phi^+ \phi^- - \frac{1}{2}\partial_\mu \phi^0 \partial_\mu \phi^0 - \frac{1}{2c_w^2}M\phi^0 \phi^0 - \beta_h[\frac{2M^2}{g^2} + \\
 & \frac{2M}{g}H + \frac{1}{2}(H^2 + \phi^0 \phi^0 + 2\phi^+ \phi^-)] + \frac{2M^4}{g^2}\alpha_h - igc_w[\partial_\nu Z_\mu^0(W_\mu^+ W_\nu^- - \\
 & W_\nu^+ W_\mu^-) - Z_\nu^0(W_\mu^+ \partial_\nu W_\mu^- - W_\nu^- \partial_\mu W_\mu^+) + Z_\mu^0(W_\nu^+ \partial_\nu W_\mu^- - \\
 & W_\nu^- \partial_\mu W_\mu^+)] - ig s_w[\partial_\nu(W_\mu^+ W_\nu^- - W_\nu^+ W_\mu^-) - A_\nu(W_\mu^+ \partial_\nu W_\mu^- - \\
 & W_\mu^- \partial_\nu W_\mu^+) + A_\mu(W_\nu^+ \partial_\nu W_\mu^- - W_\nu^- \partial_\mu W_\mu^+)] - \frac{1}{2}g^2 W_\mu^+ W_\mu^- W_\nu^+ W_\nu^- + \\
 & \frac{1}{2}g^2 W_\mu^+ W_\nu^- W_\mu^+ W_\nu^- + g^2 c_w^2(Z_\mu^0 W_\mu^+ Z_\nu^0 W_\nu^- - Z_\mu^0 Z_\nu^0 W_\mu^+ W_\nu^-) + \\
 & g^2 s_w^2(A_\mu W_\mu^+ A_\nu W_\nu^- - A_\mu A_\nu W_\mu^+ W_\nu^-) + g^2 s_w c_w[A_\mu Z_\nu^0(W_\mu^+ W_\nu^- - \\
 & W_\nu^+ W_\mu^-) - 2A_\mu Z_\mu^0 W_\nu^+ W_\nu^-] - g\alpha[H^3 + H\phi^0 \phi^0 + 2H\phi^+ \phi^-] - \\
 & \frac{1}{8}g^2 \alpha_h[H^4 + (\phi^0)^4 + 4(\phi^+ \phi^-)^2 + 4(\phi^0)^2 \phi^+ \phi^- + 4H^2 \phi^+ \phi^- + 2(\phi^0)^2 H^2] - \\
 & gMW_\mu^+ W_\mu^- H - \frac{1}{2}g\frac{M}{c_w^2}Z_\mu^0 Z_\mu^0 H - \frac{1}{2}ig[W_\mu^+(\phi^0 \partial_\mu \phi^- - \phi^- \partial_\mu \phi^0) - \\
 & W_\mu^-(\phi^0 \partial_\mu \phi^+ - \phi^+ \partial_\mu \phi^0)] + \frac{1}{2}ig[W_\mu^+(H\partial_\mu \phi^- - \phi^- \partial_\mu H) - W_\mu^-(H\partial_\mu \phi^+ - \\
 & \phi^+ \partial_\mu H)] + \frac{1}{2}ig[\frac{1}{c_w}(Z_\mu^0(H\partial_\mu \phi^0 - \phi^0 \partial_\mu H) - ig\frac{s_w^2}{c_w}M Z_\mu^0(W_\mu^+ \phi^- - W_\mu^- \phi^+) + \\
 & ig s_w A_\mu(W_\mu^+ \phi^- - W_\mu^- \phi^+) - ig\frac{1-2c_w^2}{2c_w}Z_\mu^0(\phi^+ \partial_\mu \phi^- - \phi^- \partial_\mu \phi^+) + \\
 & ig s_w A_\mu(\phi^+ \partial_\mu \phi^- - \phi^- \partial_\mu \phi^+) - \frac{1}{4}g^2 W_\mu^+ W_\mu^- [H^2 + (\phi^0)^2 + 2\phi^+ \phi^-] - \\
 & \frac{1}{4}g^2 \frac{1}{c_w^2}Z_\mu^0 Z_\mu^0 [H^2 + (\phi^0)^2 + 2(2s_w^2 - 1)^2 \phi^+ \phi^-] - \frac{1}{2}g^2 \frac{s_w^2}{c_w^2}Z_\mu^0 \phi^0(W_\mu^+ \phi^- + \\
 & W_\mu^- \phi^+) - \frac{1}{2}ig^2 \frac{s_w^2}{c_w^2}Z_\mu^0 H(W_\mu^+ \phi^- - W_\mu^- \phi^+) + \frac{1}{2}g^2 s_w A_\mu \phi^0(W_\mu^+ \phi^- + \\
 & W_\mu^- \phi^+) + \frac{1}{2}ig^2 s_w A_\mu H(W_\mu^+ \phi^- - W_\mu^- \phi^+) - g^2 \frac{s_w}{c_w}(2c_w^2 - 1)Z_\mu^0 A_\mu \phi^+ \phi^- - \\
 & g^4 s_w^4 A_\mu A_\mu \phi^+ \phi^- - \bar{e}^\lambda(\gamma^\mu \partial_\mu + m_e^\lambda)e^\lambda - \bar{\nu}^\lambda \gamma^\mu \partial_\mu \nu^\lambda - \bar{u}_j^\lambda(\gamma^\mu \partial_\mu + m_u^\lambda)u_j^\lambda - \\
 & \bar{d}_j^\lambda(\gamma^\mu \partial_\mu + m_d^\lambda)d_j^\lambda + ig s_w A_\mu [-(\bar{e}^\lambda \gamma^\mu e^\lambda) + \frac{2}{3}(\bar{u}_j^\lambda \gamma^\mu u_j^\lambda) - \frac{1}{3}(\bar{d}_j^\lambda \gamma^\mu d_j^\lambda)] + \\
 & \frac{ig}{4c_w}Z_\mu^0[(\bar{\nu}^\lambda \gamma^\mu(1 + \gamma^5)\nu^\lambda) + (\bar{e}^\lambda \gamma^\mu(4s_w^2 - 1 - \gamma^5)e^\lambda) + (\bar{u}_j^\lambda \gamma^\mu(\frac{4}{3}s_w^2 - \\
 & 1 - \gamma^5)u_j^\lambda) + (\bar{d}_j^\lambda \gamma^\mu(1 - \frac{8}{3}s_w^2 - \gamma^5)d_j^\lambda)] + \frac{ig}{2\sqrt{2}}W_\mu^+[(\bar{\nu}^\lambda \gamma^\mu(1 + \gamma^5)e^\lambda) + \\
 & (\bar{u}_j^\lambda \gamma^\mu(1 + \gamma^5)C_{\lambda\kappa}d_j^\kappa)] + \frac{ig}{2\sqrt{2}}W_\mu^-[(\bar{e}^\lambda \gamma^\mu(1 + \gamma^5)\nu^\lambda) + (\bar{d}_j^\kappa C_{\lambda\kappa}^\dagger \gamma^\mu(1 + \\
 & \gamma^5)u_j^\lambda)] + \frac{ig}{2\sqrt{2}}\frac{m_\lambda}{M}[-\phi^+(\bar{\nu}^\lambda(1 - \gamma^5)e^\lambda) + \phi^-(\bar{e}^\lambda(1 + \gamma^5)\nu^\lambda)] - \\
 & \frac{g}{2}\frac{m_\lambda}{M}[H(\bar{e}^\lambda e^\lambda) + i\phi^0(\bar{e}^\lambda \gamma^5 e^\lambda)] + \frac{ig}{2M\sqrt{2}}\phi^+[-m_d^\lambda(\bar{u}_j^\lambda C_{\lambda\kappa}(1 - \gamma^5)d_j^\kappa) + \\
 & m_u^\lambda(\bar{u}_j^\lambda C_{\lambda\kappa}(1 + \gamma^5)d_j^\kappa)] + \frac{ig}{2M\sqrt{2}}\phi^-[m_d^\lambda(\bar{d}_j^\kappa C_{\lambda\kappa}^\dagger(1 + \gamma^5)u_j^\lambda) - m_u^\lambda(\bar{d}_j^\kappa C_{\lambda\kappa}^\dagger(1 - \\
 & \gamma^5)u_j^\lambda) - \frac{g}{2}\frac{m_\lambda}{M}H(\bar{u}_j^\lambda u_j^\lambda) - \frac{g}{2}\frac{m_\lambda}{M}H(\bar{d}_j^\lambda d_j^\lambda) + \frac{ig}{2}\frac{m_\lambda}{M}\phi^0(\bar{u}_j^\lambda \gamma^5 u_j^\lambda) - \\
 & \frac{ig}{2}\frac{m_\lambda}{M}\phi^0(\bar{d}_j^\lambda \gamma^5 d_j^\lambda) + \bar{X}^+(\partial^2 - M^2)X^+ + \bar{X}^-(\partial^2 - M^2)X^- + \bar{X}^0(\partial^2 - \\
 & \frac{M^2}{c_w^2})X^0 + \bar{Y}\partial^2 Y + igc_w W_\mu^+(\partial_\mu \bar{X}^0 X^- - \partial_\mu \bar{X}^+ X^0) + ig s_w W_\mu^+(\partial_\mu \bar{Y} X^- - \\
 & \partial_\mu \bar{X}^+ Y) + igc_w W_\mu^-(\partial_\mu \bar{X}^- X^0 - \partial_\mu \bar{X}^0 X^+) + ig s_w W_\mu^-(\partial_\mu \bar{X}^- Y - \\
 & \partial_\mu \bar{Y} X^+) + igc_w Z_\mu^0(\partial_\mu \bar{X}^+ X^- - \partial_\mu \bar{X}^- X^+) + ig s_w A_\mu(\partial_\mu \bar{X}^+ X^- - \\
 & \partial_\mu \bar{X}^- X^+) - \frac{1}{2}gM[\bar{X}^+ X^+ H + \bar{X}^- X^- H + \frac{1}{c_w}\bar{X}^0 X^0 H] +
 \end{aligned}$$



Standard Model

Three generations of matter (fermions)

	I	II	III	
mass	2.4 MeV/c ²	1.27 GeV/c ²	171.2 GeV/c ²	0
charge	2/3	2/3	2/3	0
spin	1/2	1/2	1/2	1
name	u up	c charm	t top	γ photon
	4.8 MeV/c ²	104 MeV/c ²	4.2 GeV/c ²	0
	-1/3	-1/3	-1/3	0
	1/2	1/2	1/2	1
Quarks	d down	s strange	b bottom	g gluon
	<2.2 eV/c ²	<0.17 MeV/c ²	<15.5 MeV/c ²	91.2 GeV/c ²
	0	0	0	0
	1/2	1/2	1/2	1
	ν _e electron neutrino	ν _μ muon neutrino	ν _τ tau neutrino	Z ⁰ Z boson
	0.511 MeV/c ²	105.7 MeV/c ²	1.777 GeV/c ²	80.4 GeV/c ²
	-1	-1	-1	±1
Leptons	e electron	μ muon	τ tau	W [±] W boson
	1/2	1/2	1/2	1





Gauge bosons

$$\begin{aligned}
 & -\frac{1}{2}\partial_\nu g_\mu^a g_\nu^a - g_s f^{abc} \partial_\mu g_\nu^a g_\mu^b g_\nu^c - \frac{1}{4}g_s^2 f^{abc} f^{ade} g_\mu^b g_\nu^c g_\mu^d g_\nu^e + \\
 & \frac{1}{2}g_s^2(\bar{q}_i^\gamma \gamma^\mu q_j^\gamma)g_\mu^a + \bar{G}^a \partial^2 G^a + g_s f^{abc} \partial_\mu \bar{G}^a G^b g_\mu^c - \partial_\nu W_\mu^+ \partial_\nu W_\mu^- - \\
 & M^2 W_\mu^+ W_\mu^- - \frac{1}{2}\partial_\nu Z_\mu^0 \partial_\nu Z_\mu^0 - \frac{1}{2c_w^2} M^2 Z_\mu^0 Z_\mu^0 - \frac{1}{2}\partial_\mu A_\nu \partial_\mu A_\nu - \frac{1}{2}\partial_\mu H \partial_\mu H - \\
 & \frac{1}{2}m_h^2 H^2 - \partial_\mu \phi^+ \partial_\mu \phi^- - M^2 \phi^+ \phi^- - \frac{1}{2}\partial_\mu \phi^0 \partial_\mu \phi^0 - \frac{1}{2c_w^2} M \phi^0 \phi^0 - \beta_h \left[\frac{2M^2}{g^2} + \right. \\
 & \left. \frac{2M}{g} H + \frac{1}{2}(H^2 + \phi^0 \phi^0 + 2\phi^+ \phi^-) \right] + \frac{2M^4}{g^2} \alpha_h - ig_{c_w} [\partial_\nu Z_\mu^0 (W_\mu^+ W_\nu^- - \\
 & W_\nu^+ W_\mu^-) - Z_\nu^0 (W_\mu^+ \partial_\nu W_\mu^- - W_\nu^+ \partial_\mu W_\mu^-) + Z_\mu^0 (W_\nu^+ \partial_\nu W_\mu^- - \\
 & W_\nu^- \partial_\nu W_\mu^+) - ig_{s_w} [\partial_\nu (W_\mu^+ W_\nu^- - W_\nu^+ W_\mu^-) - A_\nu (W_\mu^+ \partial_\nu W_\mu^- - \\
 & W_\mu^- \partial_\nu W_\mu^+) + A_\mu (W_\nu^+ \partial_\nu W_\mu^- - W_\nu^- \partial_\nu W_\mu^+)] - \frac{1}{2}g^2 W_\mu^+ W_\mu^- W_\nu^+ W_\nu^- + \\
 & \frac{1}{2}g^2 W_\mu^+ W_\nu^- W_\mu^- W_\nu^+ + g^2 c_w^2 (Z_\mu^0 W_\mu^+ Z_\nu^0 W_\nu^- - Z_\mu^0 Z_\nu^0 W_\mu^+ W_\nu^-) + \\
 & g^2 s_w^2 (A_\mu W_\mu^+ A_\nu W_\nu^- - A_\mu A_\nu W_\mu^+ W_\nu^-) + g^2 s_w c_w [A_\mu Z_\nu^0 (W_\mu^+ W_\nu^- - \\
 & W_\nu^+ W_\mu^-) - 2A_\mu Z_\mu^0 W_\nu^+ W_\nu^-] - g\alpha [H^3 + H\phi^0 \phi^0 + 2H\phi^+ \phi^-] - \\
 & \frac{1}{8}g^2 \alpha_h [H^4 + (\phi^0)^4 + 4(\phi^+ \phi^-)^2 + 4(\phi^0)^2 \phi^+ \phi^- + 4H^2 \phi^+ \phi^- + 2(\phi^0)^2 H^2] - \\
 & g M W_\mu^+ W_\mu^- H - \frac{1}{2}g \frac{M}{c_w^2} Z_\mu^0 Z_\mu^0 H - \frac{1}{2}ig [W_\mu^+ (\phi^0 \partial_\mu \phi^- - \phi^- \partial_\mu \phi^0) - \\
 & W_\mu^- (\phi^0 \partial_\mu \phi^+ - \phi^+ \partial_\mu \phi^0)] + \frac{1}{2}ig [W_\mu^+ (H \partial_\mu \phi^- - \phi^- \partial_\mu H) - W_\mu^- (H \partial_\mu \phi^+ - \\
 & \phi^+ \partial_\mu H)] + \frac{1}{2}ig [\frac{1}{c_w} (Z_\mu^0 (H \partial_\mu \phi^0 - \phi^0 \partial_\mu H) - ig \frac{s_w^2}{c_w} M Z_\mu^0 (W_\mu^+ \phi^- - W_\mu^- \phi^+) + \\
 & ig_{s_w} M A_\mu (W_\mu^+ \phi^- - W_\mu^- \phi^+) - ig \frac{1-2c_w^2}{2c_w} Z_\mu^0 (\phi^+ \partial_\mu \phi^- - \phi^- \partial_\mu \phi^+) + \\
 & ig_{s_w} A_\mu (\phi^+ \partial_\mu \phi^- - \phi^- \partial_\mu \phi^+) - \frac{1}{4}g^2 W_\mu^+ W_\mu^- [H^2 + (\phi^0)^2 + 2\phi^+ \phi^-] - \\
 & \frac{1}{4}g^2 \frac{1}{c_w^2} Z_\mu^0 Z_\mu^0 [H^2 + (\phi^0)^2 + 2(2s_w^2 - 1)^2 \phi^+ \phi^-] - \frac{1}{2}g^2 \frac{s_w^2}{c_w^2} Z_\mu^0 \phi^0 (W_\mu^+ \phi^- + \\
 & W_\mu^- \phi^+) - \frac{1}{2}ig^2 \frac{s_w^2}{c_w} Z_\mu^0 H (W_\mu^+ \phi^- - W_\mu^- \phi^+) + \frac{1}{2}g^2 s_w A_\mu \phi^0 (W_\mu^+ \phi^- + \\
 & W_\mu^- \phi^+) + \frac{1}{2}ig^2 s_w A_\mu H (W_\mu^+ \phi^- - W_\mu^- \phi^+) - g^2 \frac{s_w}{c_w} (2c_w^2 - 1) Z_\mu^0 A_\mu \phi^+ \phi^- - \\
 & g^4 s_w^2 A_\mu A_\nu \phi^+ \phi^- - \bar{e}^\lambda (\gamma^\mu \partial_\nu + m_e^\lambda) e^\lambda - \bar{\nu}^\lambda \gamma^\mu \partial_\nu \nu^\lambda - \bar{u}_j^\lambda (\gamma^\mu \partial_\nu + m_u^\lambda) u_j^\lambda - \\
 & \bar{d}_j^\lambda (\gamma^\mu \partial_\nu + m_d^\lambda) d_j^\lambda + ig_{s_w} A_\mu [-(\bar{e}^\lambda \gamma^\mu e^\lambda) + \frac{2}{3}(\bar{u}_j^\lambda \gamma^\mu u_j^\lambda) - \frac{1}{3}(\bar{d}_j^\lambda \gamma^\mu d_j^\lambda)] + \\
 & \frac{ig}{4c_w} Z_\mu^0 [(\bar{\nu}^\lambda \gamma^\mu (1 + \gamma^5) \nu^\lambda) + (\bar{e}^\lambda \gamma^\mu (4s_w^2 - 1 - \gamma^5) e^\lambda) + (\bar{u}_j^\lambda \gamma^\mu (\frac{4}{3}s_w^2 - \\
 & 1 - \gamma^5) u_j^\lambda) + (\bar{d}_j^\lambda \gamma^\mu (1 - \frac{8}{3}s_w^2 - \gamma^5) d_j^\lambda)] + \frac{ig}{2\sqrt{2}} W_\mu^+ [(\bar{\nu}^\lambda \gamma^\mu (1 + \gamma^5) e^\lambda) + \\
 & (\bar{u}_j^\lambda \gamma^\mu (1 + \gamma^5) C_{\lambda k} d_k^\lambda)] + \frac{ig}{2\sqrt{2}} W_\mu^- [(\bar{e}^\lambda \gamma^\mu (1 + \gamma^5) \nu^\lambda) + (\bar{d}_j^\lambda C_{\lambda k}^\dagger \gamma^\mu (1 + \\
 & \gamma^5) u_j^\lambda)] + \frac{ig}{2\sqrt{2}} \frac{m_\lambda}{M} [-\phi^+ (\bar{\nu}^\lambda (1 - \gamma^5) e^\lambda) + \phi^- (\bar{e}^\lambda (1 + \gamma^5) \nu^\lambda)] - \\
 & \frac{g}{2} \frac{m_\lambda}{M} [H (\bar{e}^\lambda e^\lambda) + i\phi^0 (\bar{e}^\lambda \gamma^5 e^\lambda)] + \frac{ig}{2M\sqrt{2}} \phi^+ [-m_d^\lambda (\bar{u}_j^\lambda C_{\lambda k} (1 - \gamma^5) d_k^\lambda) + \\
 & m_u^\lambda (\bar{u}_j^\lambda C_{\lambda k} (1 + \gamma^5) d_k^\lambda) + \frac{ig}{2M\sqrt{2}} \phi^- [m_d^\lambda (\bar{d}_j^\lambda C_{\lambda k}^\dagger (1 + \gamma^5) u_j^\lambda) - m_u^\lambda (\bar{d}_j^\lambda C_{\lambda k}^\dagger (1 - \\
 & \gamma^5) u_j^\lambda) - \frac{g}{2} \frac{m_\lambda}{M} H (\bar{u}_j^\lambda u_j^\lambda) - \frac{g}{2} \frac{m_\lambda}{M} H (\bar{d}_j^\lambda d_j^\lambda) + \frac{ig}{2} \frac{m_\lambda}{M} \phi^0 (\bar{u}_j^\lambda \gamma^5 u_j^\lambda) - \\
 & \frac{ig}{2} \frac{m_\lambda}{M} \phi^0 (\bar{d}_j^\lambda \gamma^5 d_j^\lambda) + \bar{X}^+ (\partial^2 - M^2) X^+ + \bar{X}^- (\partial^2 - M^2) X^- + \bar{X}^0 (\partial^2 - \\
 & \frac{M^2}{c_w^2}) X^0 + \bar{Y} \partial^2 Y + ig_{c_w} W_\mu^+ (\partial_\mu \bar{X}^0 X^- - \partial_\mu \bar{X}^+ X^0) + ig_{s_w} W_\mu^+ (\partial_\mu \bar{Y} X^- - \\
 & \partial_\mu \bar{X}^+ Y) + ig_{c_w} W_\mu^- (\partial_\mu \bar{X}^- X^0 - \partial_\mu \bar{X}^0 X^+) + ig_{s_w} W_\mu^- (\partial_\mu \bar{X}^- Y - \\
 & \partial_\mu \bar{Y} X^+) + ig_{c_w} Z_\mu^0 (\partial_\mu \bar{X}^+ X^- - \partial_\mu \bar{X}^- X^+) + ig_{s_w} A_\mu (\partial_\mu \bar{X}^+ X^- + \\
 & \partial_\mu \bar{X}^- X^+) - \frac{1}{2}g M [\bar{X}^+ X^+ H + \bar{X}^- X^- H + \frac{1}{c_w} \bar{X}^0 X^0 H] +
 \end{aligned}$$



Experiments?




At very high energy collisions, we can probe the fundamental building blocks of nature.


We can verify whether these predictions are actually true.

We can also look for the “New Physics” in the collisions, which were proposed to solve some problems of the SM.


Experiments?




Experiments?



Large Hadron Collider




175 m underground




*Pune for size comparison

Circumference : 27 km

Experiments?




Proton




Proton

Experiments?




Proton




Proton


Experiments?




Experiments?




LHC Detectors




ATLAS/CMS Detector




ATLAS/CMS Detector



ATLAS/CMS Detector



Experiments!



Wait!



Leptonic



0.511 MeV/c²
-1
1/2 **e**
electron

105.7 MeV/c²
-1
1/2 **μ**
muon

1.777 GeV/c²
-1
1/2 **τ**
tau

Hadronic

Detector signatures of electrons & muons



**Search for new phenomena in 4 lepton final
states with the full Run 2 dataset at ATLAS
detector at LHC**

...


SHREYAS BAKARE

Search for **new phenomena** in **4 lepton final states** with the full Run 2 dataset at ATLAS detector at LHC

...


SHREYAS BAKARE

$Z \rightarrow$




Z
~~~~~


$$Z \rightarrow \tau^+ \tau^-$$



# New phenomena




# 4 lepton final state






# 4 lepton final state




# 4 lepton final state



$\mu^+ \mu^+ \tau^- \tau^- \longrightarrow$ 
 $\begin{matrix} \mu^+ \mu^+ \mu^- \mu^- & \mu^+ \mu^+ \mu^- e^- & \mu^+ \mu^+ e^- e^- \\ \mu^+ \mu^+ \mu^- h & \mu^+ \mu^+ e^- h & \mu^+ \mu^+ h h \end{matrix}$

WHY?







Calculating signal events in run 2  
for different  $Z'$  masses



# Light Lepton multiplicity (N) for $l\bar{l}$ Pt cut $>5$ GeV





So what? | Lagrangians  
to  
Lasers





So what?

**EXTRA SLIDES**

# Theoretical models: with Z boson



- **Model #1:**  $gg \rightarrow A \rightarrow ZH (\rightarrow WW/\tau\tau)$  in a 2HDM, where H can be the SM Higgs or a BSM Higgs
- **Model #2:** vector-like leptons with decay in Z boson allowed
  - Singlet and doublet models; single and pair produced (larger cross section predicted for pair produced VLL)
  - It can also accommodate  $(g-2)_{\mu/e}$  data
  - Other decays possible to a new BSM scalar **S**
  - See [2011.12964]





# Theoretical models: without Z boson (I)




- **Model #3:**  $gg \rightarrow H \rightarrow Sh/SS \rightarrow WW\tau\tau/WWWW$




- Addresses other multilepton tensions
- See [1912.00699]

- **Model #4:** leptophilic  $Z'$  or scalar with flavour-violating couplings

- Flavour off-diagonal  $Z'$  couplings to the  $\mu$  and  $\tau$  sectors [satisfies various constraints from LEP  $e^+e^- \rightarrow e^+e^-$ , the  $(g-2)_e$ , ...], *or*
- 2HDM model with sizeable couplings to  $\mu$  and  $\tau$ , leading as well to  $\mu^\pm \mu^\mp \tau^\mp \tau^\mp$  final states
- Could address the  $(g-2)_\mu$  tension
- See [1607.06832] and [1907.09845]



Preferred heavy Higgs masses  
 $\sim \mathcal{O}(100)$  GeV and limited  
 below  $\sim 700$  GeV  
 [1907.09845]





# Search for new physics at the Large Hadron Collider

...

SHREYAS BAKARE

*Experimental High Energy  
Physics*

## The details of the session

Speaker:

Shreyas Bakare

Topic:

Search for new physics at the Large Hadron Collider

Venue:


LHC 106 (6:30 PM)

Abstract:



Despite the success of the Standard Model in describing the interactions of elementary particles, observations that suggest the existence of additional phenomena remain. Many theories of physics beyond the Standard Model have been proposed that feature "final states" in high-energy proton-proton collisions with exactly four leptons. In this talk, I will start with the basics of experimental high energy physics and move towards discussing this particular 4-lepton search that targets particular mixed flavor regions. The ideas will be motivated, so there is no need for much background!

# Large Hadron Collider

CERN's Accelerator Complex





# Why not hadronic?



Detector signatures of neutral & charged hadrons

- **Model #5:** leptophilic  $Z'$  or scalar coupling preferentially to  $\tau$

- Could address the  $(g-2)_\mu$  tension
- Explore four  $\tau$  signature




Flavour off-diagonal  $Z'$  coupling  
to the  $\mu$  and  $\tau$  sectors

I Mainly worked on  $p p \rightarrow \tau \mu Z p$   
Where  $Z p$  couples to  $\tau$  &  $\mu$  (violating Lepton flavour)




# $Z'$ coupling to $\mu$ & $\tau$



- In these two plots, from Lepton flavour violating  $Z'$  explanation of the muon anomalous magnetic moment[2], the green band is preferred at  $2\sigma$  by the  $(g-2)_\mu$  anomaly, whereas the grey region is disfavored at  $> 5\sigma$ . The red region is excluded by lepton flavour universality in tau decays.
- Based on these two plots[2], we adjust the couplings as a function of  $Z'$  mass ( $m_{Z'}$ ) as

$$g_{23R} = 0.003 \text{ GeV}^{-1} \times m_{Z'} / \text{GeV}$$




$$g_{23L} = g_{23R} / 10.$$



$g_L$  v/s  $g_R$  plane for  $m_{Z'} = 100 \text{ GeV}$

$m_{Z'}$  v/s  $g_R$  for  $g_L = g_R/10$

$g-2$



# MC generation using MadGraph + Pythia

- First-ever Monte Carlo generation of the  $p p \rightarrow \tau \mu Z$  process.
- Probing Z mass in the range of 10 GeV to 100 GeV as the cross-section of the process turns out 13.76 fb for  $m_Z = 10$  GeV whereas 0.25 fb for  $m_Z = 100$  GeV.

```
p p > mu+(-) ta- (+) Zp /h NP<=2 QED<=2
```

```
1. For MZp = 10  
WZp = 5.807750e-05  
cross-section (fb)= 35.28
```


```
2. For MZp = 30  
WZp = 1.635278e-03  
cross-section (fb)= 31.21
```

```
For MZp = 50  
WZp = 7.595605e-03  
cross-section (fb)= 8.763
```


```
3. For MZp = 70  
WZp = 2.086114e-02  
cross-section (fb)= 1.132
```

```
4. For MZp = 100  
WZp = 6.084878e-02  
cross-section (fb)= 0.3798
```






# Comparing kinematics as a function of $m_Z$



# Comparing kinematics as a function of $m_{Z'}$





Flavour off-diagonal  $Z'$  coupling  
to the  $\mu$  and  $\tau$  sectors



## Generation of EVGEN.root

## Conversion into DAOD file


## Conversion into flat ntuple



WORKFLOW



# Interesting Problem



DAOD\_TRUTH1.test.pool.root

- Metadata;2
- Metadata;1
- MetadataHdr;2
- MetadataHdr;1
- MetadataHdrForm;2
- MetadataHdrForm;1
- POOLContainer;2
- POOLContainer;1
- POOLContainerForm;2
- POOLContainerForm;1
- ##Params;2
- ##Params;1
- ##Shapes;2
- ##Shapes;1
- ##Links;2
- ##Links;1
- CollectionTree;1

- TruthTausAuxDyn.eta\_vis
- TruthTausAuxDyn.phi\_vis
- TruthTausAuxDyn.m\_vis
- TruthTausAuxDyn.numCharged
- TruthTausAuxDyn.numChargedPion
- TruthTausAuxDyn.numNeutral
- TruthTausAuxDyn.numNeutralPion
- TruthTausAuxDyn.IsHadronicTau
- TruthTausAuxDyn.pt\_invis
- TruthTausAuxDyn.px
- TruthTausAuxDyn.py
- TruthTausAuxDyn.eta\_invis
- TruthTausAuxDyn.pz
- TruthTausAuxDyn.phi\_invis
- TruthTausAuxDyn.e
- TruthTausAuxDyn.m\_invis
- TruthTausAuxDyn.DecayModeVector
- TruthTausAuxDyn.pdgId
- TruthTausAuxDyn.classifierParticleType
- TruthTausAuxDyn.classifierParticleOrigin

```

//
std::vector<TLorentzVector> TruthAna_GenericSelector::HadronicTau(const xAOD::TruthParticleContainer *cont, float pt, float eta,
CUT_PDG cut_pdg, int pdgid, CUT_STATUS cut_status, int status, char IsHadronic)
{
    std::vector<TLorentzVector> tlv_vec;

    for(auto vcont : *cont){
        float loc_px    =vcont->px(); float loc_py    =vcont->py();
        float loc_pz    =vcont->pz(); float loc_e      =vcont->e() ;
        TLorentzVector tlv; tlv.SetPxPyPzE(loc_px,loc_py,loc_pz,loc_e);

        int par_pdgid=vcont->auxdata<int>("pdgId");
        int par_IsHadronic=vcont->auxdata<char>("IsHadronicTau");

        if( tlv.Pt()<pt )continue;
        if( fabs( tlv.Rapidity() )>eta )continue;

        //special treatment for the H/A bosons
        if( par_pdgid==35 || par_pdgid== 36 || fabs(par_pdgid== 37) || par_pdgid==25 ) par_pdgid=35;

        if      ( cut_pdg==CUT_PDG::YES && ( ! (par_pdgid== pdgid ) ) ) continue;
        else if( cut_pdg==CUT_PDG::ABS && ( ! (fabs(par_pdgid)== pdgid ) ) ) continue;
        if (par_IsHadronic == IsHadronic) continue;
        if( cut_status==CUT_STATUS::YES && ( ! (vcont->auxdata<int>("status")== status ) ) ) continue;

        tlv_vec.push_back(tlv);
    }

    return tlv_vec;
}
//

```

$p p \rightarrow \mu_{\pm} \tau a^{\mp} Z p$ , 4 lep final state  $Q = 0$

$$N = \epsilon \sigma L$$


Light Lepton multiplicity (N) for  $l l$  Pt cut  $> 5$  GeV

For  $M_{Zp} = 10$  GeV, 30 GeV, 50 GeV, 70 GeV & 100 GeV  
 $\sigma = 35.28$  fb, 31.21 fb, 8.763 fb, 1.132 fb & 0.3798 fb resp.  
 $L = 140$  /fb

For  $e, \mu, \tau$   $\eta < 2.5$ ,  $\tau$  pt  $> 5$  GeV




# Light Lepton multiplicity (N) for $l\bar{l}$ Pt cut $>5$ GeV



# Integrals of each histograms (w/o SS pairs condition)

| MZp | Pt > 5 GeV |
|-----|------------|
| 10  | 39901.9    |
| 30  | 59251.2    |
| 50  | 4787.79    |
| 70  | 102.036    |
| 100 | 18.3794    |

# SS pairs: Light Lepton multiplicity (N) for $l\bar{l}$ Pt cut $>5$ GeV



# Integrals of each histograms (SS pairs)

| MZp | Pt > 5 GeV |
|-----|------------|
| 10  | 24394.2    |
| 30  | 32765.6    |
| 50  | 2840.39    |
| 70  | 61.688     |
| 100 | 12.1317    |